
Polycomb-like 2 associates with PRC2 and regulates transcriptional networks during mouse embryonic stem cell self-renewal and differentiation.

Journal: Cell Stem Cell

Publication Year: 2010

Authors: Emily Walker, Wing Y Chang, Julie Hunkapiller, Gerard Cagney, Kamal Garcha, Joseph Torchia, Nevan J Krogan, Jeremy F Reiter, William L Stanford

PubMed link: 20144788

Funding Grants: High throughput modeling of human neurodegenerative diseases in embryonic stem cells

Public Summary:

We used embryonic stem cells to biochemically purify PRC2, a complex of proteins that regulates stem cell pluripotency. Analysis of this complex discovered a new regulator of pluripotency called Pcl2.

Scientific Abstract:

Polycomb group (PcG) proteins are conserved epigenetic transcriptional repressors that control numerous developmental gene expression programs and have recently been implicated in modulating embryonic stem cell (ESC) fate. We identified the PcG protein PCL2 (polycomb-like 2) in a genome-wide screen for regulators of self-renewal and pluripotency and predicted that it would play an important role in mouse ESC-fate determination. Using multiple biochemical strategies, we provide evidence that PCL2 is a Polycomb Repressive Complex 2 (PRC2)-associated protein in mouse ESCs. Knockdown of Pcl2 in ESCs resulted in heightened self-renewal characteristics, defects in differentiation, and altered patterns of histone methylation. Integration of global gene expression and promoter occupancy analyses allowed us to identify PCL2 and PRC2 transcriptional targets and draft regulatory networks. We describe the role of PCL2 in both modulating transcription of ESC self-renewal genes in undifferentiated ESCs as well as developmental regulators during early commitment and differentiation.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/polycomb-2-associates-prc2-and-regulates-transcriptional-networks-during>